

| | | | | | | | |
|--------|--------------|-----|---|------|----|----|----|
| Female | Neutered | Yes | 1 | 4.00 | 4 | 4 | 4 |
| Female | Not neutered | No | 4 | NA | NA | NA | NA |
| Male | Neutered | Yes | 1 | 2.00 | 2 | 2 | 2 |
| Male | Not neutered | No | 4 | NA | NA | NA | NA |

Table 4: The body condition comment and the age of the dog when this comment was made

| Body condition | Count | Mean age (years) | Median age (years) | Min age (years) | Max age (years) |
|---------------------|-------|------------------|--------------------|-----------------|-----------------|
| Normal | 5 | 2.80 | 2 | 1 | 5 |
| Somewhat overweight | 1 | 4.00 | 4 | 4 | 4 |

Table 5: The most commonly reported disease conditions in descending order for the breed (N = 10 live dogs)

| Condition |
|---|
| Ectropion |
| Entropion |
| Prolapsed third eyelid gland (Cherry eye) |
| Corneal ulcer |
| Corneal Ulcer |
| Dermatitis |
| Enlarged heart |
| Hypersensitivity (allergic) skin disorder |
| Irregular Heart beat |
| Osteochondritis dissecans (OCD) |
| Spondylosis |

Summary

From the dogs surveyed, most Neapolitan Mastiffs were affected by a disease condition (60.00%). There were three disease conditions that represented an equal proportion of affected live dogs; these conditions were ectropion, entropion, and prolapsed third eyelid gland (Cherry eye). The most commonly reported cause of death was gastric dilation-volvulus syndrome GDV (Bloat).